

p -adic congruences in iterated derivatives of \wp

Kiran Luecke, Eric Peterson

Abstract

We use homotopy theoretic methods to prove congruence relations of number theoretic interest. Specifically, we use the theory of \mathbb{E}_∞ complex orientations to establish p -adic K  m  er congruences among iterated derivatives of the Weierstrass elliptic function.

The machinery of [AHR] was developed by Ando, Hopkins, and Rezk with the intended application of taking congruence relations as input and producing \mathbb{E}_∞ -orientations as output. We run their machine in reverse, using as input the recent results of [CL25] on the existence of \mathbb{E}_∞ -orientations of Tate fixed-point objects.

	t^0	t^{-1}	t^{-2}	t^{-3}	t^{-4}	t^{-5}
$M_1^{\text{Wit}^\sharp}$...000000 ₂	...111110 ₂	...111110 ₂	...111100 ₂	...111000 ₂	...110000 ₂
$M_2^{\text{Wit}^\sharp}$...101010 ₂	...111000 ₂	...111000 ₂	...100000 ₂	...110000 ₂	...000000 ₂
$M_3^{\text{Wit}^\sharp}$...000000 ₂	...100110 ₂	...100110 ₂	...111100 ₂	...101000 ₂	...100000 ₂
$M_4^{\text{Wit}^\sharp}$...011010 ₂	...110000 ₂	...110000 ₂	...000000 ₂	...100000 ₂	...000000 ₂
$M_5^{\text{Wit}^\sharp}$...000000 ₂	...001110 ₂	...001110 ₂	...111100 ₂	...011000 ₂	...010000 ₂
$M_6^{\text{Wit}^\sharp}$...001010 ₂	...101000 ₂	...101000 ₂	...100000 ₂	...010000 ₂	...000000 ₂
$M_7^{\text{Wit}^\sharp}$...000000 ₂	...110110 ₂	...110110 ₂	...111100 ₂	...001000 ₂	...000000 ₂
$M_8^{\text{Wit}^\sharp}$...111010 ₂	...100000 ₂	...100000 ₂	...000000 ₂	...000000 ₂	...000000 ₂
$M_9^{\text{Wit}^\sharp}$...000000 ₂	...011110 ₂	...011110 ₂	...111100 ₂	...111000 ₂	...110000 ₂
$M_{10}^{\text{Wit}^\sharp}$...101010 ₂	...011000 ₂	...011000 ₂	...100000 ₂	...110000 ₂	...000000 ₂
$M_{11}^{\text{Wit}^\sharp}$...000000 ₂	...000110 ₂	...000110 ₂	...111100 ₂	...101000 ₂	...100000 ₂
$M_{12}^{\text{Wit}^\sharp}$...011010 ₂	...010000 ₂	...010000 ₂	...000000 ₂	...100000 ₂	...000000 ₂
$M_{13}^{\text{Wit}^\sharp}$...000000 ₂	...101110 ₂	...101110 ₂	...111100 ₂	...011000 ₂	...010000 ₂
$M_{14}^{\text{Wit}^\sharp}$...001010 ₂	...001000 ₂	...001000 ₂	...100000 ₂	...010000 ₂	...000000 ₂
$M_{15}^{\text{Wit}^\sharp}$...000000 ₂	...010110 ₂	...010110 ₂	...111100 ₂	...001000 ₂	...000000 ₂
$M_{16}^{\text{Wit}^\sharp}$...111010 ₂	...000000 ₂				
$M_{17}^{\text{Wit}^\sharp}$...000000 ₂	...111110 ₂	...111110 ₂	...111100 ₂	...111000 ₂	...110000 ₂
$M_{18}^{\text{Wit}^\sharp}$...101010 ₂	...111000 ₂	...111000 ₂	...100000 ₂	...110000 ₂	...000000 ₂
$M_{19}^{\text{Wit}^\sharp}$...000000 ₂	...100110 ₂	...100110 ₂	...111100 ₂	...101000 ₂	...100000 ₂
$M_{20}^{\text{Wit}^\sharp}$...011010 ₂	...110000 ₂	...110000 ₂	...000000 ₂	...100000 ₂	...000000 ₂
$M_{21}^{\text{Wit}^\sharp}$...000000 ₂	...001110 ₂	...001110 ₂	...111100 ₂	...011000 ₂	...010000 ₂
$M_{22}^{\text{Wit}^\sharp}$...001010 ₂	...101000 ₂	...101000 ₂	...100000 ₂	...010000 ₂	...000000 ₂
$M_{23}^{\text{Wit}^\sharp}$...000000 ₂	...110110 ₂	...110110 ₂	...111100 ₂	...001000 ₂	...000000 ₂
$M_{24}^{\text{Wit}^\sharp}$...111010 ₂	...100000 ₂	...100000 ₂	...000000 ₂	...000000 ₂	...000000 ₂
$M_{25}^{\text{Wit}^\sharp}$...000000 ₂	...011110 ₂	...011110 ₂	...111100 ₂	...111000 ₂	...110000 ₂
$M_{26}^{\text{Wit}^\sharp}$...101010 ₂	...011000 ₂	...011000 ₂	...100000 ₂	...110000 ₂	...000000 ₂
$M_{27}^{\text{Wit}^\sharp}$...000000 ₂	...000110 ₂	...000110 ₂	...111100 ₂	...101000 ₂	...100000 ₂
$M_{28}^{\text{Wit}^\sharp}$...011010 ₂	...010000 ₂	...010000 ₂	...000000 ₂	...100000 ₂	...000000 ₂
$M_{29}^{\text{Wit}^\sharp}$...000000 ₂	...101110 ₂	...101110 ₂	...111100 ₂	...011000 ₂	...010000 ₂
$M_{30}^{\text{Wit}^\sharp}$...001010 ₂	...001000 ₂	...001000 ₂	...100000 ₂	...010000 ₂	...000000 ₂
$M_{31}^{\text{Wit}^\sharp}$...000000 ₂	...010110 ₂	...010110 ₂	...111100 ₂	...001000 ₂	...000000 ₂

Contents

1	Introduction	2
2	Résumés	3
2.1	Résumé on elliptic functions	3
2.2	Résumé on elliptic cohomology	5
2.3	Résumé on \mathbb{E}_∞ -rings	7
3	Proofs of the Main Theorems	9
3.1	Todd and Todd $^\sharp$	9
3.2	Witten and Witten $^\sharp$	10
4	Loose ends	11

1 Introduction

Definition 1.1 ([Kat77, Sections 3.1–2]). For A a torsion-free \mathbb{Z}_p -algebra, an A -valued p -adic moment sequence is a sequence $M_n \in A$ such that for any rational polynomial $f(r) = \sum_n a_n r^n \in \mathbb{Q}[r]$ with $f(\mathbb{Z}_p^\times) \subseteq \mathbb{Z}_p$, the sum $\sum_n a_n M_n \in A \otimes \mathbb{Q}$ is an A -integer.

Moment sequences arise out of p -adic integrals via the correspondence

$$\int_{\mathbb{Z}_p^\times} dM \sum_{n=0}^N a_n r^n := \sum_{n=0}^N a_n M_n, \quad M_n := \int_{\mathbb{Z}_p^\times} dM r^n.$$

The integrality condition can be interpreted as a family of *Kümmel congruences* on the sequence elements. For example, the polynomial $p^{-1}(r^p - r)$ is p -adically integer-valued, hence $p^{-1}(M_p - M_1)$ must be an A -integer, or equivalently $M_p \equiv M_1 \pmod{p}$. Other choices of test polynomial yield other congruences.

We prove the following:

Main Theorem A. *Pick any $c \in \mathbb{Z}_p^\times \setminus \{\pm 1\}$. There is a $\mathbb{Z}[[q]]((t))_p^\wedge$ -valued moment sequence whose terms for $n \geq 1$ are given by*

$$\begin{aligned} M_1^{\text{Wit}^\sharp} &= (1 - c) \cdot \frac{\log(1 - t)}{(2\pi i)^2} \cdot (G_2(q) - p \cdot G_2(q^p)) \\ &\quad + \frac{1 - c}{2\pi i} \cdot \left(\wp_q^{(-1)} \left(\frac{\log(1 - t)}{-2\pi i} \right) - \wp_{q^p}^{(-1)} \left(\frac{p \log(1 - t)}{-2\pi i} \right) \right), \\ M_2^{\text{Wit}^\sharp} &= (1 - c^2) \cdot \frac{G_2(q) - p \cdot G_2(q^p)}{-2(2\pi i)^2} \\ &\quad + (1 - c^2) \cdot \left(\wp_q \left(\frac{\log(1 - t)}{-2\pi i} \right) - \wp_{q^p} \left(\frac{p \log(1 - t)}{-2\pi i} \right) \right), \\ M_{n \geq 3}^{\text{Wit}^\sharp} &= \frac{1 - c^n}{(2\pi i)^n} \cdot \left(\wp_q^{(n-2)} \left(\frac{\log(1 - t)}{-2\pi i} \right) - \wp_{q^p}^{(n-2)} \left(\frac{p \log(1 - t)}{-2\pi i} \right) \right), \end{aligned}$$

where \wp_q is the q -expansion of the Weierstrass elliptic function, $\wp_q^{(n)}$ is its n^{th} derivative, and G_2 is the second Eisenstein series.

As warm-up, we also prove the following theorem:

Main Theorem B. *Pick any $c \in \mathbb{Z}_p^\times \setminus \{\pm 1\}$. There is a $\mathbb{Z}((t))_p^\wedge$ -valued moment sequence whose terms for $n \geq 1$ are given by*

$$(1 - c^n) \left(\sum_{m=1}^n \frac{\Delta^m[r^n](0)}{m} ((t^{-1})^m - p^{n-1}t^{-pm}(1 - p\zeta)^{-m}) \right),$$

where $\Delta^m[r^n](0)$ is the evaluation at zero of m^{th} forward finite difference of the test polynomial r^n , and where $\zeta \in \mathbb{Z}[t^{-1}]$ is defined by $(t^{-1} - 1)^p = t^{-p} - 1 + p\zeta$.

Remark 1.2. The p in front of ζ yields a convergent sum in $\mathbb{Z}((t))_p^\wedge$, despite the unboundedly increasing powers of t^{-1} .

Remark 1.3. Main Theorem B is essentially the component of Main Theorem A induced by the pole of $\wp^{(n-2)}$. More exactly:

- The two Main Theorems' terms of nonpositive t -degree match exactly.
- The two Main Theorems' terms of vanishing t - and q -degree are exactly the moments of the Mazur measure [Lan90, Section 4.3].
- The terms of Main Theorem A of vanishing t -degree are exactly those of the $MF_{*,*}$ -valued measure constructed by Katz [Kat77, Lemma 3.5.6], appearing also in Ando, Hopkins, and Rezk [AHR, Section 10].

Remark 1.4. See Figure 1 and Figure 2 for visualizations of these congruences at $p = 3$, and Figure 3 and Figure 4 (which extends the title figure) for $p = 2$.

Acknowledgments

We would like to thank Charles Rezk for helpful conversations, Jonathan Beardsley for helpful comments on a draft, and Matthew Ando, Shachar Carmeli, and Tomer Schlank for their words of encouragement. The second author would like to thank BART, where much of this work was carried out.

2 Résumés

Our main theorems are largely an application of existing work, especially a recent paper of the first author [CL25]. Accordingly, we recall below the inputs which we require to complete our calculation.

2.1 Résumé on elliptic functions

It will be convenient to have formulas on hand for a variety of standard functions appearing in the theory of complex elliptic curves. To fix terms, let us consider (twisted) functions on the compactified moduli of elliptic curves, referred to as *modular forms*. Such functions are typically presented in three ways [Sil94, Section I.1, Remark 3.3]:

1. A modular form f of weight k is a function on choices of lattice $\Lambda \subset \mathbb{C}$ with the property $f(c \cdot \Lambda) = c^{-k} f(\Lambda)$ and which extends to a meromorphic function at $\text{span}\{+i \cdot \infty, 1\}$.
2. Selecting a basis $\{\tau, 1\}$ for the (possibly positively rescaled) lattice Λ , f can be understood as a function of $\tau \in \mathfrak{h}$ satisfying $f(M \cdot \tau) = (c\tau + d)^{-k} f(\tau)$ and extending to a meromorphic function at $\tau = +i \cdot \infty$.
3. Using periodicity against $\text{SL}_2\mathbb{Z}$ and meromorphicity at $\tau = +i \cdot \infty$, we may write the Fourier series of f as a bounded-below sum $\sum_{n=-m}^{\infty} a_n q^n$, where $q = e^{2\pi i \tau}$, called the q -expansion.¹

We will also be interested in functions on the universal elliptic curve, called *Jacobi forms* [EZ85, Section I.1], modeled as functions on $\mathbb{C} \times \mathfrak{h}$ equivariant for $\text{SL}_2\mathbb{Z} \ltimes \mathbb{Z}^2$. Such functions again can be parameterized in terms of Λ , τ , or q .

Example 2.1. For $k \geq 2$, the k^{th} Eisenstein series G_k is the (quasi²)modular form defined by the lattice sum [Kat77, Equation 2.2.5]

$$G_k(\Lambda) = \sum_{\lambda \in \Lambda^*} \frac{1}{\lambda^k}.$$

It is evidently of weight k . Note that if k is odd, the sum telescopes to zero. In the even case, the q -expansion of G_{2k} is

$$\frac{(2k-1)!}{(2\pi i)^{2k}} \cdot G_{2k}(q) = -\frac{B_{2k}}{2k} + 2 \cdot \sum_{j=1}^{\infty} \sigma_{2k-1}(j) q^j,$$

where B_{2k} is the $2k^{\text{th}}$ Bernoulli number, defined by $\sum_{n=0}^{\infty} B_n x^n = \frac{x}{e^x - 1}$, and where $\sigma_{2k-1}(j) = \sum_{d|j} d^{2k-1}$ is the divisor-sum function [Kat77, Equation 2.4.8].

Example 2.2. The Weierstrass σ -function is defined by [Sil94, Proposition 5.4.a]

$$\sigma_{\Lambda}(z) = z \cdot \prod_{\lambda \in \Lambda^*} \left[\left(1 - \frac{z}{\lambda}\right) \cdot \exp \left(\frac{z}{\lambda} + \frac{1}{2} \left(\frac{z}{\lambda} \right)^2 \right) \right].$$

Among other properties, it is designed so that its logarithm becomes a generating function for Eisenstein series [Kat77, Equation 2.2.7]:

$$\log z - \log \sigma_{\Lambda}(z) = \sum_{n=3}^{\infty} \frac{(2\pi iz)^n}{n!} \cdot \left(\frac{(n-1)!}{(2\pi i)^n} \cdot G_n(\Lambda) \right).$$

We will need a product formula for its q -expansion [Sil94, Theorems 6.4, 6.3.b]:

$$2\pi i \cdot \sigma_q(z) = \exp \left(\frac{1}{2} (2\pi i \cdot z + G_2(q) \cdot z^2) \right) \cdot (1-u) \cdot \prod_{j=1}^{\infty} \frac{(1-q^j u)(1-q^j u^{-1})}{(1-q^j)^2},$$

¹However, it is not trivial to discern when an arbitrary q -series is a modular form.

² $G_2(\tau)$ does not quite transform as a modular form, but instead acquires an extra summand.

where $u = e^{-2\pi iz}$. Lastly, the second logarithmic derivative of σ is named the *Weierstrass \wp -function* [Sil94, Proposition 5.4.c]:

$$\wp_\Lambda(z) = -\frac{d^2}{dz^2} \log \sigma_\Lambda(z),$$

which is a meromorphic Jacobi form of index 0 and weight 2.

2.2 Résumé on elliptic cohomology

In this subsection we recall some basic elements of complex orientation theory.

Definition 2.3. Let R be a homotopy ring spectrum. The following two pieces of data are equivalent:

1. A *complex orientation* of R is a homotopy ring map $\mathrm{MU} \rightarrow R$ from the complex bordism spectrum [Ada74, Lemma I.4.6].
2. A *coordinate* is a class $\xi \in R^2\mathbb{CP}^\infty$ which restricts to $\Sigma^2 1 \in R^2 S^2$ along $S^2 \simeq \mathbb{CP}^1 \rightarrow \mathbb{CP}^\infty$ [Ada74, I.2.1].

A coordinate induces an isomorphism $R_*[\![\xi]\!] \rightarrow R^* \mathbb{CP}^\infty$. The Künneth formula then gives an isomorphism $R^*(\mathbb{CP}^\infty \times \mathbb{CP}^\infty) \simeq R_*[\![\xi_l, \xi_r]\!]$ where ξ_l and ξ_r are the pullbacks of ξ along the two projections. Let $\mu: \mathbb{CP}^\infty \times \mathbb{CP}^\infty \rightarrow \mathbb{CP}^\infty$ be the multiplication map classifying the tensor product of line bundles. The *formal group law* $F(\xi_l, \xi_r)$ is defined to be the power series associated to the element $\mu^* \xi$ [Ada74, Lemma I.2.7]. We often denote it by $\xi_l +_F \xi_r$.

Example 2.4. Note that MU is a connective ring spectrum with $\pi_0 \mathrm{MU} = \mathbb{Z}$, so the Postnikov truncation functor induces a ring map $\mathrm{MU} \rightarrow \mathbb{Z}$. Thus, if R is a \mathbb{Z} -algebra, there is a canonical orientation $\mathrm{MU} \rightarrow \mathbb{Z} \rightarrow R$. We will refer to this as the *additive* orientation of R , as its group law is $F(x_l, x_r) = x_l + x_r$.

Example 2.5. Write $\pi_* \mathrm{KU} = \mathbb{Z}[\beta^\pm]$ and $\pi_* \mathrm{KU}_{\mathrm{Tate}} = \mathbb{Z}[\beta^\pm][[q]]$ with $|\beta| = 2$ and $|q| = 0$.

- The *Todd orientation* ω_{Td} : $\mathrm{MU} \rightarrow \mathrm{KU}$ has coordinate [Ada74, I.2.3]

$$\xi_{\mathrm{Td}} := \beta^{-1}(1 - [L]) \in \mathrm{KU}^2 \mathbb{CP}^\infty.$$

- The *Witten orientation* ω_{Wit} : $\mathrm{MU} \rightarrow \mathrm{KU}_{\mathrm{Tate}}$ has coordinate [Zag88, (6)]

$$\xi_{\mathrm{Wit}} = \beta^{-1}(1 - [L]) \prod_{j=1}^{\infty} \frac{(1 - [L]q^j)(1 - [L^{-1}]q^j)}{(1 - q^j)^2} \in \mathrm{KU}_{\mathrm{Tate}}^2 \mathbb{CP}^\infty.$$

For later convenience we also define the degree zero element $s := \beta \xi_{\mathrm{Td}}$.

Definition 2.6 ([Ada74, Corollary I.7.15]). Let $\omega: \mathrm{MU} \rightarrow R$ be an orientation, which begets an orientation $\omega_{\mathbb{Q}}: \mathrm{MU} \rightarrow R \otimes \mathbb{Q}$. Since $R \otimes \mathbb{Q}$ is a \mathbb{Z} -algebra, it has an additive orientation (cf. Example 2.4) with coordinate x . Write ξ_{ω} for the coordinate associated to $\omega_{\mathbb{Q}}$. In terms of x , it is a power series of the form $\exp_{\omega}(x) := \xi_{\omega}(x) = x + O(x^2)$. It is called the *exponential* of the formal group law F_{ω} associated to ω and satisfies the formula

$$\exp_{\omega}(x_l) +_{F_{\omega}} \exp_{\omega}(x_r) = \exp_{\omega}(x_l + x_r).$$

Example 2.7 ([Hir66, Chapter 3]). The exponential of the Todd orientation is

$$\exp_{\mathrm{Td}}(x) = \beta^{-1}(1 - e^{-\beta x}).$$

Therefore we have

$$\exp_{\mathrm{Wit}}(x) = \beta^{-1}(1 - e^{-\beta x}) \prod_{j=1}^{\infty} \frac{(1 - e^{-\beta x} q^j)(1 - e^{\beta x} q^j)}{(1 - q^j)^2}.$$

Definition 2.8 ([AHR, Section 3.4]). The *nepers* of an orientation $\omega: \mathrm{MU} \rightarrow R$ are the elements $N_n^{\omega} \in \pi_{2n}R \otimes \mathbb{Q}$ defined by the formula

$$\log\left(\frac{x}{\exp_{\omega}(x)}\right) = \sum_{n=1}^{\infty} N_n^{\omega} \frac{x^n}{n!}$$

When R has a distinguished periodicity element $\beta \in \pi_2R$, it will often be convenient to work with the degree zero element $\beta^{-n}N_n^{\omega}$ instead of N_n^{ω} .

Remark 2.9 ([AHR, Proposition 3.12]). The division of the two Thom classes for the universal line bundle $L \rightarrow \mathbb{CP}^{\infty}$ is classically referred to as the *Hirzebruch series* of ω . One can form the same division of Thom classes for any vector bundle, and the neper N_n^{ω} is what one obtains for the bundle over S^n corresponding to a generator of $\pi_{2n}\mathrm{BU}$.

Next, we outline a construction for getting new orientations from old ones, due to Ando, French, and Ganter [AFG08]. First, we recall a fundamental piece of homotopy theory.

Definition 2.10 (e.g., [GM95], [NS18]). Let R be a complex oriented ring spectrum with coordinate γ . The *Tate fixed-point spectrum* $R^{t\mathbb{T}}$ is the ring spectrum obtained by inverting the element γ in the ring spectrum $R^{\mathbb{CP}^{\infty}}$. In particular $\pi_*R^{t\mathbb{T}}$ is the Laurent series ring $R_*((\gamma))$, and there is a *unit* map $u: R \rightarrow R^{t\mathbb{T}}$ of ring spectra, inducing the obvious inclusion $R_* \rightarrow R_*((\gamma))$ on homotopy groups.

Definition 2.11 ([AFG08]). Let $\omega: \mathrm{MU} \rightarrow R$ be a complex orientation with its induced isomorphism $\pi_*R^{t\mathbb{T}} \simeq R_*((\gamma))$. Define the *unit orientation* of $R^{t\mathbb{T}}$ as the composite

$$\mathrm{MU} \xrightarrow{\omega} R \xrightarrow{u} R^{t\mathbb{T}},$$

and let ξ and $+_F$ denote the corresponding coordinate and formal group law. Define the *sharped orientation* $\omega^\sharp: \mathrm{MU} \rightarrow R^{t\mathbb{T}}$ to be the ring map associated to the coordinate

$$\xi \cdot \frac{\gamma}{\xi +_F \gamma}.$$

Let y be such that $\exp_\omega(y) = \gamma$. Then the sharped exponential satisfies

$$\exp_{\omega^\sharp}(x) = \exp_\omega(x) \cdot \frac{\exp_\omega(y)}{\exp_\omega(x+y)}.$$

Remark 2.12 ([AFG08, Sections 6–7]). The *Jacobi orientation* arises as the sharp construction applied to the Witten orientation. On homotopy, it takes values in meromorphic Jacobi forms of index 0.

2.3 Résumé on \mathbb{E}_∞ -rings

Whereas in classical algebra commutativity is a *property* of a ring and maps of rings are automatically compatible with any commutativity present, a ring spectrum in homotopy theory requires *extra data* to witness its commutativity, as do maps of commutative ring spectra. A ring spectrum with such extra structure is called an \mathbb{E}_∞ -ring, and such a map an \mathbb{E}_∞ -ring map. The complex bordism spectrum MU is naturally an \mathbb{E}_∞ -ring ([May77, Chapter III]), and thus if R is an \mathbb{E}_∞ -ring one can ask when a complex orientation $\mathrm{MU} \rightarrow R$ admits an \mathbb{E}_∞ -structure. This question has received much interest and has seen recent progress ([AHR], [HL18], [HY20], [Bal23], [BSY24], …), but it is usually quite difficult—the number of known \mathbb{E}_∞ -orientations is small, the crown jewel of which is the *string orientation* of tmf.

Our interest comes from a result of the first author:

Theorem 2.13 ([CL25, Theorem 5.14]). *The sharp construction sends \mathbb{E}_∞ -orientations to \mathbb{E}_∞ -orientations.* \square

Therefore the sharped string orientation (or any of the handful of \mathbb{E}_∞ -orientations it begets, cf. Theorem 2.20) automatically admits this extra \mathbb{E}_∞ -structure, and we would like to understand it in detail. Ultimately, we will conclude that it entails the congruences advertised in the introduction.

Let us describe others' pursuit of the program to express this \mathbb{E}_∞ -structure in a manageable form for palatable target rings. There is a distinguished class of spectra, called *$K(1)$ -local*, whose \mathbb{E}_∞ -rings carry an extremely nice theory of cohomology operations, controlled by an essentially unique operation ψ^p [BMMS86, Chapter IX]. A key insight of Ando, Hopkins, and Rezk is that if R is $K(1)$ -local, then an \mathbb{E}_∞ -structure on an orientation $\omega: \mathrm{MU} \rightarrow R$ is well captured by its interaction with ψ^p . Our rings of interest, $\mathrm{KU}^{t\mathbb{T}}$ and $\mathrm{KU}_{\mathrm{Tate}}^{t\mathbb{T}}$, become $K(1)$ -local after p -completion. Their ψ^p maps are as follows:

Example 2.14. Upon p -completion, the Todd and Witten orientations of Example 2.5 induce identifications

$$\pi_*(\mathrm{KU}^{t\mathbb{T}})_p^\wedge = \mathbb{Z}[\beta^\pm]((t))_p^\wedge, \quad \pi_*(\mathrm{KU}_{\mathrm{Tate}}^{t\mathbb{T}})_p^\wedge = \mathbb{Z}[\beta^\pm][[q]]((t))_p^\wedge,$$

where we have adjusted the Laurent element $t := \beta\gamma$ into degree 0 (cf. Definition 2.10). The operations ψ^p are the ring homomorphisms determined by

$$\psi^p(t) = 1 - (1-t)^p, \quad \psi^p(q) = q^p, \quad \psi^p(\beta) = p\beta.$$

For a homotopy element $f_q(t) \cdot \beta^k \in \pi_{2k} \mathrm{KU}_{\mathrm{Tate}}^{t\mathbb{T}} \cong \mathbb{Z}[[q]]((t))_p^\wedge \{\beta^k\}$ we then have

$$\psi^p(f_q(t) \cdot \beta^k) = p^k f_{q^p}(1 - (1-t)^p) \cdot \beta^k.$$

We measure the interaction of ψ^p with the orientation ω like so:

Definition 2.15 ([AHR, Proposition 7.10]). Pick any $c \in \mathbb{Z}_p^\times \setminus \{\pm 1\}$. The *moments* of an orientation $\omega: \mathrm{MU} \rightarrow R$ of a $K(1)$ -local ring are the elements

$$M_n^\omega = (1 - c^n) \left(\mathrm{id} - \frac{\psi^p}{p} \right) (N_n^\omega) \in \pi_{2n} R \otimes \mathbb{Q}.$$

As at the end of Definition 2.8, when R has a distinguished periodicity element $\beta \in \pi_2 R$, it will often be convenient to work with the degree 0 element $\beta^{-n} M_n^\omega$. However, one has to take a bit of care, since $\psi^p(\beta)$ need not be equal to β (as Example 2.14 shows).

Remark 2.16 ([Rez06]). The operator $\mathrm{id} - p^{-1}\psi^p$ is part of a more general theory of *logarithmic cohomology operations*.

Remark 2.17 ([AHR, Proposition 14.6]). The factor $1 - c^n$ comes from an *Adams resolution* of the $K(1)$ -localized sphere spectrum.

We can now state a key result of Ando–Hopkins–Rezk:

Theorem 2.18 ([AHR, Theorem 6.1], [Pet19, Appendix A.4]). Let $\omega: \mathrm{MU} \rightarrow R$ be an \mathbb{E}_∞ -orientation of a $K(1)$ -local ring R which is flat over KU_p^\wedge . The moments $M_{n \geq 1}^\omega$ belong to a p -adic moment sequence with values in R_* . \square

Remark 2.19 ([AHR, Lemma 7.14, Proposition 14.6.2, Section 3.5], [Mil82]). Orientations are automatically suitably compatible with the unit map of the ring spectrum, which manifests in the moment sequence as the identity

$$\lim_{j \rightarrow \infty} M_{(p-1)p^j} = M_0 = p^{-1} \log c^{p-1}.$$

Finally, we need some existing \mathbb{E}_∞ -orientations of KU and $\mathrm{KU}_{\mathrm{Tate}}$ to which we will apply Theorem 2.13 to get \mathbb{E}_∞ -orientations of $(\mathrm{KU}^{t\mathbb{T}})_p^\wedge$ and $(\mathrm{KU}_{\mathrm{Tate}}^{t\mathbb{T}})_p^\wedge$, and then further apply Theorem 2.18 to get p -adic moment sequences. We record the following theorem.

Theorem 2.20 ([AHR, Theorem 10.3, Proposition 10.10]). After p -completion, the orientations of Example 2.5

$$\omega_{\mathrm{Td}}: \mathrm{MU} \rightarrow \mathrm{KU}_p^\wedge, \quad \omega_{\mathrm{Wit}}: \mathrm{MU} \rightarrow (\mathrm{KU}_{\mathrm{Tate}})_p^\wedge$$

admit \mathbb{E}_∞ -structures. \square

Remark 2.21. To be precise, the statement of Proposition 10.10 of [AHR] produces an \mathbb{E}_∞ -structure on the $MU\langle 6 \rangle$ -orientation associated to ω_{Wit} . However, the same proof, with the target ring changed to π_*KU_{Tate} , shows that ω_{Wit} itself admits an \mathbb{E}_∞ -orientation. One can also use obstruction theory along the Whitehead tower of bu to push the \mathbb{E}_∞ -structure from $MU\langle 6 \rangle$ to MU .

3 Proofs of the Main Theorems

We now apply this machinery to the sharped orientations of (Tate) K -theory from Theorem 2.20 to produce p -adic moment sequences. It will be convenient to fix the following notation:

$$2\pi iz = x = -\log(1-s) = -\log(u), \quad 2\pi i\alpha = y = -\log(1-t) = -\log(v).$$

We remark that while s agrees with the element defined at the end of Example 2.5, x does not quite agree with the additive coordinate of previous sections—it is the degree 0 element gotten by using the periodicity element β . This slight abuse of notation is well worth it, as the formulas that follow would otherwise be infested with powers of β .

Throughout, let $c \in \mathbb{Z}_p^\times \setminus \{\pm 1\}$ be a choice of p -adic unit.

3.1 Todd and $Todd^\sharp$

Lemma 3.1 ([Kob84, Theorem 7.2], [AHR, Proposition 10.2]). *There is a p -adic moment sequence which in positions $n \geq 1$ is given by*

$$(1 - c^n)(1 - p^{n-1}) \frac{-B_n}{n},$$

where B_n denotes the n^{th} Bernoulli number.

Proof. Using Example 2.7, the nepers associated to the Todd orientation are

$$\sum_{n=0}^{\infty} N_n^{\text{Td}} \frac{x^n}{n!} = \log\left(\frac{x}{s}\right) = \log\left(\frac{x}{1 - e^{-x}}\right).$$

The series expansion of the right-hand side follows from:

$$\frac{d}{dx} \log \frac{x}{1 - e^{-x}} = \frac{1}{x} - \frac{e^{-x}}{1 - e^{-x}} = \frac{1}{x} - \frac{1}{x} \cdot \frac{x}{e^x - 1} = \sum_{n=1}^{\infty} -B_n \frac{x^{n-1}}{n!}.$$

We then calculate the moments (cf. Definition 2.15) and find that they are the claimed sequence, which is a p -adic moment sequence by Theorem 2.18. \square

Proof of Main Theorem B. Combining Example 2.7 and Definition 2.11, the nepers associated to the $Todd^\sharp$ orientation are

$$\sum_{n=0}^{\infty} N_n^{\text{Td}^\sharp} \frac{x^n}{n!} = \log\left(\frac{x}{s}\right) + \log\left(\frac{1 - e^{-x}(1-t)}{t}\right).$$

The first summand is the subject of Lemma 3.1, so we focus on the second:

$$\log \left(\frac{1 - t^{-1}(1 - e^x)}{e^x} \right) = -x - \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{m=1}^n \frac{t^{-m}}{m} \sum_{j=0}^m (-1)^j \binom{m}{j} j^n.$$

The inner sum is exactly the iterated forward finite difference $\Delta^m[r^n](0)$. Apply Theorem 2.18 to produce the p -adic moment sequence:

$$M_n^{\text{Td}\#} = (1 - c^n) \left(\sum_{m=1}^n \frac{\Delta^m[r^n](0)}{m} (t^{-m} - p^{n-1}(1 - (1-t)^p)^{-m}) \right).$$

The binomial raised to the $(-m)^{\text{th}}$ power does not naively expand, as $(1-t)^p$ is not a topologically nilpotent element of $\pi_*(K^{t\mathbb{T}})_p^\wedge$. However, by defining ζ such that $(t^{-1} - 1)^p = t^{-p} - 1 + p\zeta$, we find:

$$M_n^{\text{Td}\#} = (1 - c^n) \left(\sum_{m=1}^n \frac{\Delta^m[r^n](0)}{m} ((t^{-1})^m - p^{n-1}t^{-pm}(1 - p\zeta)^{-m}) \right),$$

which does define a convergent series in $\pi_*(K^{t\mathbb{T}})_p^\wedge$. \square

Remark 3.2. The t^{-m} term in $N_n^{\text{Td}\#}$ vanishes when $m > n$ for two reasons: the m -fold forward finite difference operator annihilates r^n , and the term $(1 - e^{-x})^m$ appearing in the expansion of $\log(1 - t^{-1}(1 - e^x))$ has x -degree at least m .

3.2 Witten and Witten $^\sharp$

Lemma 3.3 ([Kat77, Lemma 3.5.6], [AHR, Propositions 10.9–10]). *There is a $\mathbb{Z}[[q]]_p^\wedge$ -valued moment sequence which in even positions $2n \geq 1$ is given by*

$$\frac{(1 - c^{2n}) \cdot (2n - 1)!}{(2\pi i)^{2n}} \cdot (G_{2n}(q) - p^{2n-1}G_{2n}(q^p))$$

and which vanishes in odd positions $2n + 1 \geq 1$.

Proof. The exponential for the Witten orientation of Tate K -theory in Example 2.7 shows the associated sequence of nepers to be

$$\sum_{n=1}^{\infty} N_n^{\text{Wit}} \frac{x^n}{n!} = \log \left(\frac{x}{1-u} \right) + \log \left(\prod_{j=1}^{\infty} \frac{(1-q^j)^2}{(1-uq^j)(1-u^{-1}q^j)} \right).$$

The first summand is felled by Lemma 3.1, so we concentrate on the second:

$$\begin{aligned} \log \left(\prod_{j=1}^{\infty} \frac{(1-q^j)^2}{(1-uq^j)(1-u^{-1}q^j)} \right) &= \sum_{j=1}^{\infty} \sum_{d=1}^{\infty} \frac{1}{d} (-2q^{jd} + q^{jd}e^{-dx} + q^{jd}e^{dx}) \\ &= 2 \sum_{n=1}^{\infty} \frac{x^{2n}}{(2n)!} \sum_{j=1}^{\infty} q^j \sigma_{2n-1}(j). \end{aligned}$$

Comparing with Example 2.1 shows

$$N_n^{\text{Wit}} = \frac{(2n-1)!}{(2\pi i)^{2n}} G_{2n}(q).$$

All that remains is to calculate the moments M_n^{Wit} and apply Theorem 2.18. \square

Proof of Main Theorem A. Again, combining Example 2.7 and Definition 2.11, the sharped Witten nepers are given by

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} N_n^{\text{Wit}\#} = \log\left(\frac{x}{\exp_{\text{Wit}}(x)}\right) + \log\left(\frac{\exp_{\text{Wit}}(x+y)}{\exp_{\text{Wit}}(y)}\right).$$

The first summand is felled by Lemma 3.3, so we focus on expanding the second. Note from Example 2.2 that the Witten exponential can be written in terms of σ_q as

$$\exp_{\text{Wit}}(x) = 2\pi i \cdot \sigma_q(z) \cdot \exp\left(-\frac{1}{2}(2\pi i \cdot z + G_2(q) \cdot z^2)\right),$$

and hence that second summand can be written as

$$\begin{aligned} \log\left(\frac{\exp_{\text{Wit}}(x+y)}{\exp_{\text{Wit}}(y)}\right) &= \left(-\frac{1}{2} + \frac{G_2(q)}{(2\pi i)^2} \log(1-t)\right) \cdot x + \frac{-G_2(q)}{2(2\pi i)^2} \cdot x^2 \\ &\quad + \sum_{n=1}^{\infty} \frac{x^n}{n!} \cdot \frac{1}{(2\pi i)^n} \cdot \wp_q^{(n-2)}\left(\frac{\log(1-t)}{-2\pi i}\right). \end{aligned}$$

The moments are then calculated to be exactly the expressions appearing in the Theorem statement, so Theorem 2.18 finishes the proof. \square

4 Loose ends

Remark 4.1. The recipe followed here applies to any other \mathbb{E}_∞ -orientations the reader may have on hand.

- [AHR]: The spectrum TMF is a topological incarnation of the moduli stack of elliptic curves over a general base. Its homotopy, analogous to the cohomology of the moduli stack, brings to light certain torsion phenomena not visible in the homotopy of $\pi_* \text{KU}_{\text{Tate}}$. The σ -orientation of tmf refines the Witten orientation of KU_{Tate} and is itself \mathbb{E}_∞ , suggesting one possible starting point for a generalization of our results.
- [HL16]: Topologists have also constructed variations on TMF for other congruence subgroups (though lifts of the σ -orientation have only been considered in limited situations). The number theory literature makes substantial use of these other stacks, and so an enrichment of our results to that setting would provide much better contact.

- [LY25]: Topologists have recently constructed a spectrum TJF which refines the theory of Jacobi forms analogously to how TMF refines the moduli of elliptic curves. This perhaps forms a more appropriate target for parametric Jacobi orientations than $\mathrm{TMF}^{t\mathbb{T}}$. (We thank Yamashita for bringing this to our attention.)
- [MT91, Appendix II]: Mazur and Tate construct functions analogous to the classical Weierstrass function, but over other adic rings and specified by quite different formulas. Since TMF lies over a general base, the σ -orientation suitably restricted may shed light on these other σ -functions.
- [Bal23], [BSY24]: Morava E -theories are known to carry \mathbb{E}_∞ -orientations, and the relevant logarithms bear a number-theoretic description [Rez06, 1.10]. To our knowledge, the associated congruences have not been explored—nevermind their sharper versions.

Remark 4.2. It is not clear to the authors how to unify the many derivatives appearing in this document. For example, it is tempting to interpret the finite differences appearing in Main Theorem B as belonging to a kind of pairing:

$$\left\langle \log\left(\frac{x}{\exp_{\mathrm{Td}^\sharp}(x)}\right), x^n t^{-m} \right\rangle = \frac{1}{n!m} \Delta^m[r^n](0),$$

where the role of t^{-m} is to select the iterated finite difference operator $\Delta^m|_{r=0}$ and x^n to select the test polynomial r^n . This interpretation does not obviously extend to the positive t -terms appearing in $N_n^{\mathrm{Wit}^\sharp}$, but *some* total interpretation seems surely possible.

Remark 4.3 ([Sof97, Section 3]). Sofer explores a construction which extracts from a Jacobi form a p -adic measure valued in modular forms, using formulas of similar flavor to ours but “with one fewer variable”. It could be interesting to have some identification of the *measure* in Main Theorem A in those terms, or to generalize their construction to produce measures valued in Jacobi forms, or even to see the limitations of any such generalizations.

More concretely, it would be interesting to see topology-free proofs of the congruences described in our Main Theorems. One specific opportunity for improvement is that Ando, Hopkins, and Rezk’s manual analysis allows them to “halve” the Mazur measure [AHR, Section 10.3]. Figure 3 and Figure 4 suggest that this is possible for the measure constructed in Main Theorem A; perhaps this is accessible through topology, but it is likely equally accessible directly.

Remark 4.4. The three factors appearing in Lemma 3.1 have discernable proveances in both homotopy theory and in number theory. In homotopy theory, they belong to the finite Adams resolution of the $K(1)$ -local sphere, the Rezk logarithm, and the characteristic series of the Todd orientation. In number theory, they arise from the regularization of a distribution to a measure, the restriction to \mathbb{Z}_p^\times to perform p -adic interpolation, and as special values of the ζ -function. It would be deeply interesting to have a tighter correspondence between these three pieces. The Main Theorems also provide a new proving ground for any

such analogy, which should additionally assign a measure-theoretic meaning to $\psi^p(t) = 1 - (1-t)^p = [p]_{\widehat{\mathbb{G}_m}}(t)$.

Remark 4.5. In its incarnation as an operation on \mathbb{E}_∞ -rings (cf. [CL25]), the sharp construction is part of a more general theory of \mathbb{E}_∞ -orientations of Tate objects. For example, this theory also subsumes the *Frobenius homomorphism* of [NS18]. One may wonder what the moment sequence associated to the Frobenius twist of the Witten orientation is.

References

- [Ada74] J. F. Adams. *Stable homotopy and generalised homology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, Ill.-London, 1974.
- [AFG08] Matthew Ando, Christopher P. French, and Nora Ganter. The Jacobi orientation and the two-variable elliptic genus. *Algebr. Geom. Topol.*, 8(1):493–539, 2008.
- [AHR] Matthew Ando, Michael J. Hopkins, and Charles Rezk. Multiplicative orientations of ko -theory and the spectrum of topological modular forms. <https://rezk.web.illinois.edu/koandtmf.pdf>.
- [Bal23] William Balderrama. Algebraic theories of power operations. *J. Topol.*, 16(4):1543–1640, 2023.
- [BMMS86] R. R. Bruner, J. P. May, J. E. McClure, and M. Steinberger. *H_∞ ring spectra and their applications*, volume 1176 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986.
- [BSY24] R. Burkland, T. Schlank, and A. Yuan. The chromatic nullstellensatz. *Ann. Math.*, 2024.
- [CL25] Shachar Carmeli and Kiran Luecke. Tate-valued characteristic classes, 2025, 2503.12134.
- [EZ85] Martin Eichler and Don Zagier. *The theory of Jacobi forms*, volume 55 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1985.
- [GM95] J. P. C. Greenlees and J. P. May. Generalized Tate cohomology. *Mem. Amer. Math. Soc.*, 113(543):viii+178, 1995.
- [Hir66] F. Hirzebruch. *Topological methods in algebraic geometry*, volume Band 131 of *Die Grundlehren der mathematischen Wissenschaften*. Springer-Verlag New York, Inc., New York, enlarged edition, 1966.
- [HL16] Michael Hill and Tyler Lawson. Topological modular forms with level structure. *Invent. Math.*, 203(2):359–416, 2016.

- [HL18] Michael J. Hopkins and Tyler Lawson. Strictly commutative complex orientation theory. *Math. Z.*, 290(1-2):83–101, 2018.
- [HY20] Jeremy Hahn and Allen Yuan. Exotic multiplications on periodic complex bordism. *J. Topol.*, 13(4):1839–1852, 2020.
- [Kat77] Nicholas M. Katz. The Eisenstein measure and p -adic interpolation. *Amer. J. Math.*, 99(2):238–311, 1977.
- [Kob84] Neal Koblitz. *p -adic numbers, p -adic analysis, and zeta-functions*, volume 58 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1984.
- [Lan90] Serge Lang. *Cyclotomic fields I and II*, volume 121 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1990. With an appendix by Karl Rubin.
- [LY25] Ying-Hsuan Lin and Mayuko Yamashita. Topological Elliptic Genera I – The mathematical foundation, 2025, 2412.02298.
- [May77] J. Peter May. *E_∞ ring spaces and E_∞ ring spectra*, volume Vol. 577 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1977. With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave.
- [Mil82] Haynes Miller. Universal Bernoulli numbers and the S^1 -transfer. In *Current trends in algebraic topology, Part 2 (London, Ont., 1981)*, volume 2 of *CMS Conf. Proc.*, pages 437–449. Amer. Math. Soc., Providence, RI, 1982.
- [MT91] B. Mazur and J. Tate. The p -adic sigma function. *Duke Math. J.*, 62(3):663–688, 1991.
- [NS18] Thomas Nikolaus and Peter Scholze. On topological cyclic homology. *Acta Math.*, 221(2):203–409, 2018.
- [Pet19] Eric Peterson. *Formal geometry and bordism operations*, volume 177 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2019.
- [Rez06] Charles Rezk. The units of a ring spectrum and a logarithmic cohomology operation. *J. Amer. Math. Soc.*, 19(4):969–1014, 2006.
- [Sil94] Joseph H. Silverman. *Advanced topics in the arithmetic of elliptic curves*, volume 151 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994.
- [Sof97] Adriana Sofr. p -adic aspects of Jacobi forms. *J. Number Theory*, 63(2):191–202, 1997.

- [Zag88] Don Zagier. Note on the Landweber-Stong elliptic genus. In *Elliptic curves and modular forms in algebraic topology (Princeton, NJ, 1986)*, volume 1326 of *Lecture Notes in Math.*, pages 216–224. Springer, Berlin, 1988.

	t^0	t^{-1}	t^{-2}	t^{-3}	t^{-4}	t^{-5}	t^{-6}
$M_1^{\text{Td}\#}$... 0000 ₃	... 2220 ₃	... 0000 ₃	... 0010 ₃	... 0100 ₃	... 0200 ₃	... 1000 ₃
$M_2^{\text{Td}\#}$... 1102 ₃	... 2110 ₃	... 0120 ₃	... 1200 ₃	... 2000 ₃	... 1000 ₃	... 1100 ₃
$M_3^{\text{Td}\#}$... 0000 ₃	... 0200 ₃	... 1000 ₃	... 1100 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_4^{\text{Td}\#}$... 1022 ₃	... 2120 ₃	... 0010 ₃	... 0200 ₃	... 2200 ₃	... 0000 ₃	... 0000 ₃
$M_5^{\text{Td}\#}$... 0000 ₃	... 1010 ₃	... 1100 ₃	... 1120 ₃	... 2100 ₃	... 2200 ₃	... 0000 ₃
$M_6^{\text{Td}\#}$... 2212 ₃	... 1100 ₃	... 0200 ₃	... 0000 ₃	... 2000 ₃	... 0000 ₃	... 2000 ₃
$M_7^{\text{Td}\#}$... 0000 ₃	... 2020 ₃	... 1000 ₃	... 2210 ₃	... 1100 ₃	... 2200 ₃	... 1000 ₃
$M_8^{\text{Td}\#}$... 2002 ₃	... 2210 ₃	... 1020 ₃	... 2200 ₃	... 0000 ₃	... 1000 ₃	... 1100 ₃
$M_9^{\text{Td}\#}$... 0000 ₃	... 2000 ₃	... 0000 ₃	... 1000 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{10}^{\text{Td}\#}$... 0222 ₃	... 1220 ₃	... 0210 ₃	... 2200 ₃	... 2200 ₃	... 0000 ₃	... 1000 ₃
$M_{11}^{\text{Td}\#}$... 0000 ₃	... 1110 ₃	... 1100 ₃	... 1020 ₃	... 1100 ₃	... 0200 ₃	... 0000 ₃
$M_{12}^{\text{Td}\#}$... 1112 ₃	... 2200 ₃	... 1100 ₃	... 0000 ₃	... 1000 ₃	... 0000 ₃	... 1000 ₃
$M_{13}^{\text{Td}\#}$... 0000 ₃	... 1120 ₃	... 2000 ₃	... 2110 ₃	... 2100 ₃	... 1200 ₃	... 1000 ₃
$M_{14}^{\text{Td}\#}$... 2202 ₃	... 0010 ₃	... 1220 ₃	... 0200 ₃	... 1000 ₃	... 1000 ₃	... 1100 ₃
$M_{15}^{\text{Td}\#}$... 0000 ₃	... 0100 ₃	... 2000 ₃	... 0200 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{16}^{\text{Td}\#}$... 0122 ₃	... 1020 ₃	... 1110 ₃	... 1200 ₃	... 2200 ₃	... 0000 ₃	... 2000 ₃
$M_{17}^{\text{Td}\#}$... 0000 ₃	... 1210 ₃	... 1100 ₃	... 0220 ₃	... 0100 ₃	... 1200 ₃	... 0000 ₃
$M_{18}^{\text{Td}\#}$... 0012 ₃	... 1000 ₃	... 2000 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{19}^{\text{Td}\#}$... 0000 ₃	... 0220 ₃	... 0000 ₃	... 2010 ₃	... 0100 ₃	... 0200 ₃	... 1000 ₃
$M_{20}^{\text{Td}\#}$... 0102 ₃	... 0110 ₃	... 2120 ₃	... 1200 ₃	... 2000 ₃	... 1000 ₃	... 1100 ₃
$M_{21}^{\text{Td}\#}$... 0000 ₃	... 1200 ₃	... 1000 ₃	... 0100 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{22}^{\text{Td}\#}$... 0022 ₃	... 0120 ₃	... 2010 ₃	... 0200 ₃	... 2200 ₃	... 0000 ₃	... 0000 ₃
$M_{23}^{\text{Td}\#}$... 0000 ₃	... 2010 ₃	... 1100 ₃	... 0120 ₃	... 2100 ₃	... 2200 ₃	... 0000 ₃
$M_{24}^{\text{Td}\#}$... 1212 ₃	... 2100 ₃	... 2200 ₃	... 0000 ₃	... 2000 ₃	... 0000 ₃	... 2000 ₃
$M_{25}^{\text{Td}\#}$... 0000 ₃	... 0020 ₃	... 1000 ₃	... 1210 ₃	... 1100 ₃	... 2200 ₃	... 1000 ₃
$M_{26}^{\text{Td}\#}$... 1002 ₃	... 0210 ₃	... 0020 ₃	... 2200 ₃	... 0000 ₃	... 1000 ₃	... 1100 ₃
$M_{27}^{\text{Td}\#}$... 0000 ₃						
$M_{28}^{\text{Td}\#}$... 2222 ₃	... 2220 ₃	... 2210 ₃	... 2200 ₃	... 2200 ₃	... 0000 ₃	... 1000 ₃
$M_{29}^{\text{Td}\#}$... 0000 ₃	... 2110 ₃	... 1100 ₃	... 0020 ₃	... 1100 ₃	... 0200 ₃	... 0000 ₃
$M_{30}^{\text{Td}\#}$... 0112 ₃	... 0200 ₃	... 0100 ₃	... 0000 ₃	... 1000 ₃	... 0000 ₃	... 1000 ₃
$M_{31}^{\text{Td}\#}$... 0000 ₃	... 2120 ₃	... 2000 ₃	... 1110 ₃	... 2100 ₃	... 1200 ₃	... 1000 ₃
$M_{32}^{\text{Td}\#}$... 1202 ₃	... 1010 ₃	... 0220 ₃	... 0200 ₃	... 1000 ₃	... 1000 ₃	... 1100 ₃
$M_{33}^{\text{Td}\#}$... 0000 ₃	... 1100 ₃	... 2000 ₃	... 2200 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{34}^{\text{Td}\#}$... 2122 ₃	... 2020 ₃	... 0110 ₃	... 1200 ₃	... 2200 ₃	... 0000 ₃	... 2000 ₃
$M_{35}^{\text{Td}\#}$... 0000 ₃	... 2210 ₃	... 1100 ₃	... 2220 ₃	... 0100 ₃	... 1200 ₃	... 0000 ₃
$M_{36}^{\text{Td}\#}$... 2012 ₃	... 2000 ₃	... 1000 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃	... 0000 ₃
$M_{37}^{\text{Td}\#}$... 0000 ₃	... 1220 ₃	... 0000 ₃	... 1010 ₃	... 0100 ₃	... 0200 ₃	... 1000 ₃
$M_{38}^{\text{Td}\#}$... 2102 ₃	... 1110 ₃	... 1120 ₃	... 1200 ₃	... 2000 ₃	... 1000 ₃	... 1100 ₃

Figure 1: A range of ternary expansions of the 3-adic Todd[#] moments at $c = 1+p$. Main Theorem B with test polynomial $p^{-i}(r^{(p-1)p^i} - 1)$ shows that the i^{th} digit within a column is periodic with periodicity $(p-1)p^i$. The colors are there to help the eye catch the vertical patterns.

$M_1^{\text{Wit}^\sharp}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...2220 ₃	...2210 ₃				
q^2	...2010 ₃	...2020 ₃	...2100 ₃	...2110 ₃	...2200 ₃	...2100 ₃
q^3	...2220 ₃	...2210 ₃				
q^4	...1210 ₃	...2000 ₃	...1120 ₃	...2210 ₃	...2020 ₃	...1110 ₃
q^5	...0120 ₃	...0210 ₃	...1200 ₃	...0220 ₃	...2100 ₃	...1200 ₃
q^6	...2010 ₃	...2020 ₃	...2100 ₃	...2110 ₃	...2200 ₃	...2100 ₃
q^7	...0000 ₃	...0020 ₃	...1110 ₃	...1200 ₃	...2010 ₃	...1020 ₃
q^8	...0020 ₃	...0210 ₃	...2200 ₃	...1020 ₃	...1100 ₃	...2200 ₃
q^9	...2220 ₃	...2210 ₃				

$M_7^{\text{Wit}^\sharp}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...2020 ₃	...1110 ₃				
q^2	...1210 ₃	...0120 ₃	...2100 ₃	...1010 ₃	...1200 ₃	...0100 ₃
q^3	...2020 ₃	...1110 ₃				
q^4	...2110 ₃	...1000 ₃	...0220 ₃	...0110 ₃	...2120 ₃	...2010 ₃
q^5	...0220 ₃	...0110 ₃	...1200 ₃	...0020 ₃	...0100 ₃	...0200 ₃
q^6	...1210 ₃	...0120 ₃	...2100 ₃	...1010 ₃	...1200 ₃	...0100 ₃
q^7	...1000 ₃	...1120 ₃	...0010 ₃	...0200 ₃	...1210 ₃	...0120 ₃
q^8	...0120 ₃	...1110 ₃	...2200 ₃	...1120 ₃	...2100 ₃	...1200 ₃
q^9	...2020 ₃	...1110 ₃				

$M_{19}^{\text{Wit}^\sharp}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...0220 ₃	...1210 ₃				
q^2	...1010 ₃	...0020 ₃	...2100 ₃	...1110 ₃	...2200 ₃	...2100 ₃
q^3	...0220 ₃	...1210 ₃				
q^4	...0210 ₃	...2000 ₃	...2120 ₃	...1210 ₃	...0020 ₃	...0110 ₃
q^5	...1120 ₃	...2210 ₃	...1200 ₃	...1220 ₃	...2100 ₃	...1200 ₃
q^6	...1010 ₃	...0020 ₃	...2100 ₃	...1110 ₃	...2200 ₃	...2100 ₃
q^7	...0000 ₃	...1020 ₃	...0110 ₃	...1200 ₃	...1010 ₃	...2020 ₃
q^8	...1020 ₃	...2210 ₃	...2200 ₃	...2020 ₃	...1100 ₃	...2200 ₃
q^9	...0220 ₃	...1210 ₃				

Figure 2: Some 3-adic Witten $^\sharp$ moments at $c = 1 + p$, in a range of t - and q -degrees. Congruences here relate the colored digits across different tables rather than within columns. The relevant test polynomials are $p^{-2}(r^{(p-1)p} - 1)$ and $p^{-3}(r^{(p-1)p^2} - 1)$.

	t^0	t^{-1}	t^{-2}	t^{-3}	t^{-4}	t^{-5}
$M_1^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1111110_2$	$\cdots 1111110_2$	$\cdots 1111100_2$	$\cdots 1111000_2$	$\cdots 1110000_2$
$M_2^{\text{Td}^\sharp}$	$\cdots 0101010_2$	$\cdots 1111000_2$	$\cdots 1111000_2$	$\cdots 1100000_2$	$\cdots 0110000_2$	$\cdots 1000000_2$
$M_3^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1100110_2$	$\cdots 1100110_2$	$\cdots 1111100_2$	$\cdots 0101000_2$	$\cdots 1100000_2$
$M_4^{\text{Td}^\sharp}$	$\cdots 1011010_2$	$\cdots 0110000_2$	$\cdots 0110000_2$	$\cdots 1000000_2$	$\cdots 1100000_2$	$\cdots 0000000_2$
$M_5^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0001110_2$	$\cdots 0001110_2$	$\cdots 1111100_2$	$\cdots 1011000_2$	$\cdots 1010000_2$
$M_6^{\text{Td}^\sharp}$	$\cdots 0001010_2$	$\cdots 0101000_2$	$\cdots 0101000_2$	$\cdots 0100000_2$	$\cdots 0010000_2$	$\cdots 1000000_2$
$M_7^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1110110_2$	$\cdots 1110110_2$	$\cdots 1111100_2$	$\cdots 0001000_2$	$\cdots 1000000_2$
$M_8^{\text{Td}^\sharp}$	$\cdots 0111010_2$	$\cdots 1100000_2$	$\cdots 1100000_2$	$\cdots 0000000_2$	$\cdots 1000000_2$	$\cdots 0000000_2$
$M_9^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0011110_2$	$\cdots 0011110_2$	$\cdots 1111100_2$	$\cdots 0111000_2$	$\cdots 0110000_2$
$M_{10}^{\text{Td}^\sharp}$	$\cdots 1101010_2$	$\cdots 1011000_2$	$\cdots 1011000_2$	$\cdots 1100000_2$	$\cdots 1110000_2$	$\cdots 1000000_2$
$M_{11}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0000110_2$	$\cdots 0000110_2$	$\cdots 1111100_2$	$\cdots 1101000_2$	$\cdots 0100000_2$
$M_{12}^{\text{Td}^\sharp}$	$\cdots 0011010_2$	$\cdots 0010000_2$	$\cdots 0010000_2$	$\cdots 1000000_2$	$\cdots 0100000_2$	$\cdots 0000000_2$
$M_{13}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0101110_2$	$\cdots 0101110_2$	$\cdots 1111100_2$	$\cdots 0011000_2$	$\cdots 0010000_2$
$M_{14}^{\text{Td}^\sharp}$	$\cdots 1001010_2$	$\cdots 0001000_2$	$\cdots 0001000_2$	$\cdots 0100000_2$	$\cdots 1010000_2$	$\cdots 1000000_2$
$M_{15}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0010110_2$	$\cdots 0010110_2$	$\cdots 1111100_2$	$\cdots 1001000_2$	$\cdots 0000000_2$
$M_{16}^{\text{Td}^\sharp}$	$\cdots 1111010_2$	$\cdots 1000000_2$	$\cdots 1000000_2$	$\cdots 0000000_2$	$\cdots 0000000_2$	$\cdots 0000000_2$
$M_{17}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0111110_2$	$\cdots 0111110_2$	$\cdots 1111100_2$	$\cdots 1111000_2$	$\cdots 1110000_2$
$M_{18}^{\text{Td}^\sharp}$	$\cdots 0101010_2$	$\cdots 0111000_2$	$\cdots 0111000_2$	$\cdots 1100000_2$	$\cdots 0110000_2$	$\cdots 1000000_2$
$M_{19}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0100110_2$	$\cdots 0100110_2$	$\cdots 1111100_2$	$\cdots 0101000_2$	$\cdots 1100000_2$
$M_{20}^{\text{Td}^\sharp}$	$\cdots 1011010_2$	$\cdots 1110000_2$	$\cdots 1110000_2$	$\cdots 1000000_2$	$\cdots 1100000_2$	$\cdots 0000000_2$
$M_{21}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1001110_2$	$\cdots 1001110_2$	$\cdots 1111100_2$	$\cdots 1011000_2$	$\cdots 1010000_2$
$M_{22}^{\text{Td}^\sharp}$	$\cdots 0001010_2$	$\cdots 1101000_2$	$\cdots 1101000_2$	$\cdots 0100000_2$	$\cdots 0010000_2$	$\cdots 1000000_2$
$M_{23}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 0110110_2$	$\cdots 0110110_2$	$\cdots 1111100_2$	$\cdots 0001000_2$	$\cdots 1000000_2$
$M_{24}^{\text{Td}^\sharp}$	$\cdots 0111010_2$	$\cdots 0100000_2$	$\cdots 0100000_2$	$\cdots 0000000_2$	$\cdots 1000000_2$	$\cdots 0000000_2$
$M_{25}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1011110_2$	$\cdots 1011110_2$	$\cdots 1111100_2$	$\cdots 0111000_2$	$\cdots 0110000_2$
$M_{26}^{\text{Td}^\sharp}$	$\cdots 1101010_2$	$\cdots 0011000_2$	$\cdots 0011000_2$	$\cdots 1100000_2$	$\cdots 1110000_2$	$\cdots 1000000_2$
$M_{27}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1000110_2$	$\cdots 1000110_2$	$\cdots 1111100_2$	$\cdots 1101000_2$	$\cdots 0100000_2$
$M_{28}^{\text{Td}^\sharp}$	$\cdots 0011010_2$	$\cdots 1010000_2$	$\cdots 1010000_2$	$\cdots 1000000_2$	$\cdots 0100000_2$	$\cdots 0000000_2$
$M_{29}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1101110_2$	$\cdots 1101110_2$	$\cdots 1111100_2$	$\cdots 0011000_2$	$\cdots 0010000_2$
$M_{30}^{\text{Td}^\sharp}$	$\cdots 1001010_2$	$\cdots 1001000_2$	$\cdots 1001000_2$	$\cdots 0100000_2$	$\cdots 1010000_2$	$\cdots 1000000_2$
$M_{31}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1010110_2$	$\cdots 1010110_2$	$\cdots 1111100_2$	$\cdots 1001000_2$	$\cdots 0000000_2$
$M_{32}^{\text{Td}^\sharp}$	$\cdots 1111010_2$	$\cdots 0000000_2$				
$M_{33}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1111110_2$	$\cdots 1111110_2$	$\cdots 1111100_2$	$\cdots 1111000_2$	$\cdots 1110000_2$
$M_{34}^{\text{Td}^\sharp}$	$\cdots 0101010_2$	$\cdots 1111000_2$	$\cdots 1111000_2$	$\cdots 1100000_2$	$\cdots 0110000_2$	$\cdots 1000000_2$
$M_{35}^{\text{Td}^\sharp}$	$\cdots 0000000_2$	$\cdots 1100110_2$	$\cdots 1100110_2$	$\cdots 1111100_2$	$\cdots 0101000_2$	$\cdots 1100000_2$
$M_{36}^{\text{Td}^\sharp}$	$\cdots 1011010_2$	$\cdots 0110000_2$	$\cdots 0110000_2$	$\cdots 1000000_2$	$\cdots 1100000_2$	$\cdots 0000000_2$

Figure 3: A range of binary expansions of the 2-adic Todd $^\sharp$ moments at $c = 1 + p$. Main Theorem B with test polynomial $2^{-(i+2)}(r^{2^i} - 1)$ shows that the 0th, 1st, and 2nd digit columns are 2-periodic, and then after those the i^{th} digit column is 2^{i-1} -periodic.

$M_1^{\text{Wit}^\#}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...111110 ₂	...111100 ₂				
q^2	...111110 ₂	...111100 ₂				
q^3	...000110 ₂	...010100 ₂	...100000 ₂	...101000 ₂	...111000 ₂	...110000 ₂
q^4	...111110 ₂	...111100 ₂				
q^5	...011010 ₂	...000000 ₂	...111100 ₂	...100100 ₂	...110100 ₂	...101000 ₂
q^6	...000110 ₂	...010100 ₂	...100000 ₂	...101000 ₂	...111000 ₂	...110000 ₂

$M_3^{\text{Wit}^\#}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...100110 ₂	...001100 ₂				
q^2	...100110 ₂	...001100 ₂				
q^3	...001110 ₂	...110100 ₂	...110000 ₂	...011000 ₂	...101000 ₂	...010000 ₂
q^4	...100110 ₂	...001100 ₂				
q^5	...110010 ₂	...110000 ₂	...011100 ₂	...100100 ₂	...110100 ₂	...101000 ₂
q^6	...001110 ₂	...110100 ₂	...110000 ₂	...011000 ₂	...101000 ₂	...010000 ₂

$M_5^{\text{Wit}^\#}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...001110 ₂	...011100 ₂				
q^2	...001110 ₂	...011100 ₂				
q^3	...010110 ₂	...010100 ₂	...000000 ₂	...001000 ₂	...011000 ₂	...110000 ₂
q^4	...001110 ₂	...011100 ₂				
q^5	...001010 ₂	...100000 ₂	...111100 ₂	...100100 ₂	...110100 ₂	...101000 ₂
q^6	...010110 ₂	...010100 ₂	...000000 ₂	...001000 ₂	...011000 ₂	...110000 ₂

$M_9^{\text{Wit}^\#}$	t^6	t^5	t^4	t^3	t^2	t^1
q^1	...011110 ₂	...111100 ₂				
q^2	...011110 ₂	...111100 ₂				
q^3	...100110 ₂	...010100 ₂	...100000 ₂	...101000 ₂	...111000 ₂	...110000 ₂
q^4	...011110 ₂	...111100 ₂				
q^5	...111010 ₂	...000000 ₂	...111100 ₂	...100100 ₂	...110100 ₂	...101000 ₂
q^6	...100110 ₂	...010100 ₂	...100000 ₂	...101000 ₂	...111000 ₂	...110000 ₂

Figure 4: Some 2-adic Witten $^\#$ moments at $c = 1 + p$, in a range of t - and q -degrees. Congruences here relate the colored digits across different tables rather than within columns. Congruences here relate the colored digits across different tables rather than within columns. The relevant test polynomials are $p^{-3}(r^2 - 1)$, $p^{-4}(r^4 - 1)$, and $p^{-5}(r^8 - 1)$.